

25-19-9
AGE
40 DOUBLE PUNCH

MALE SEX F

DIED DURING OR AFTER ANAES.
NONE
OPERATIVE OR OTHER
ANAESTHETIC RESPONSIBLE
MALAISE VOMITING
OTHER
MECHANICAL RESPIRATOR
MANUAL ASS. OR C.R.
T. PIECE
SEMI OPEN
ABSORPTION
ENDOTRACHEAL
ENDBRONCHIAL ANAES. OR BLOCKER
NONE
OTHER
ETHER
C₂H₆
T.C.E.
HAZELANE
N₂O
NON-DEPOLARIZER
DEPOLARIZER
NONE
OTHER
HYPOTENSIVE
PETHIDINE OR OPIATE
BARBITURATE

POST-OP. COMPS. 3

INHALATIONAL TECHNIQUES *Nalox*

INHALATIONAL AGENTS

INTRAVENOUS DRUGS

RECORD OF SPECIAL INTEREST

EMERGENCY
GOOD
FAIR
POOR
CIRCULATORY DECOMPENSATION
SEVERE RESP. DISTURBANCE
VOMITING OR FULL STOMACH
OTHER ANAES. HANDICAP
ATROPINE OR SCOPOLAMINE
PETHIDINE OR OPIATE
PHENYLETHYLAMINE DERIVATIVE
BARBITURATE
OTHER

PRE-OP. COMPLICATIONS & CONDITION
PREMEDICATION

DATE 1-6-72 No. 133

ANAESTHETIC RECORD

ANAESTHETIST *FRB*

NAME *Aegum Aesolome.* AGE *30*
HOSP. & WARD *Salalah Civil* WT. *50kg*
B.P. *140/-* T.P.R. Hb.

SURGEON *Cetti.*
OPERATION PERFORMED *D.P.S.*

PREOPERATIVE FINDINGS :-

Debondement 27.5.72
Chest clear.
Trachea central.

POSTOPERATIVE PROGRESS :-

PREMEDICATION :-

FINAL RESULT :-

SITE OF OPERATION

DISTURBANCES DURING ANAES. & THERAPY

REGIONAL ANALGESIA

2 1 1/2
3 HRS. ANAES.

SPINE & BACK
INTRA-CRANIAL
MOUTH, NOSE & THROAT
THYROID
OTHER HEAD & NECK
HEART & GT. VESSELS
PULMONARY
OTHER INTRA-THORACIC
CHEST WALL
LUMBAR
UPPER ABDOMINAL
LOWER ABDOMINAL
INGUINAL
PERINEAL
EXTREMITIES

NO OP. : DIAG. :
EXP. : THERAP. :
CARD. ARREST
CONVULSION
EXPLOSION
OTHER
CIRCULATORY
UNCOMPENSATED RESPIRATORY
UNPROTECTED VOMITING
OTHER
NONE
BLOOD TRANSFUSION
OTHER
I.V. FLUID
EXTRA-CORP. CIRCULATION
HYPOTHERMIA
OTHER

SPINAL
EPIDURAL
OTHER
NONE

PARAMOUNT REGD. TRADE MARK REGD./C.-8177 B. PRINTED IN ENGLAND.

INDIVIDUAL PRIVATE CODE


A B C D E F G H J K L M N O P R S T U

CODE: ANAESTHESIA: OP. O. C.: B.P. V.: PULSE: %: RESP. O.:
 CHART EACH DOSE AS GIVEN

TIME 1050 1100 1110 1120 1130 1140

VOL. %
 BARB. 1.
 2.
 3.
 4.
 5.

50 - 0.6 250



PATIENT'S NAME & No.
Aegum Aesalome.

| DRUGS & DOSAGE | |
|----------------|--------|
| BARBITURATE | % |
| 1. Thio | 25 250 |
| 2. Atrop. | 0.6. |
| 3. Prew | 50 |
| 4. | |
| 5. | |

COTT. 8-0 xs
 Movementful.

ANAESTHESIST FieB
 PREMED. & EFFECT — √+
None

TEETH: CHEST MOVT. GOOD POOR INVISIBLE
 PRE-OP. VEINS: GOOD POOR INVISIBLE
 SITE OF BARB. INJ. R.O ARM WRIST HAND
 ANKLE FOOT
 SITE OF I.V. FLUIDS R.L. ARM WRIST HAND
2L ANKLE FOOT
 ARMBOARD CANULA POLYTHENE CUT-DOWN
 PHAR. PRO NASOPHARYNGEAL CUP PACK
 BLIND R.L. ENDOBRONCH. BUCKETS TRACHY.
 ART. VENTILATION: INDUCTION / MAINTENANCE
 POSTURE TILT EFFECT TIME

POSTOPERATIVE STATE: Satisfactory WEAK WAKE RESTLESS PH. REFLEX.
 C.V.S. DEPRESSION RESP. DEPRESSION/DISTRESS/OBSTRUCTION VOMIT

TIME OF NOTES

